



Jay and Dani,
Real Life
Livmarli Peers



Connect with a LIVMARLI Peer Today

You are not alone!

Whether you or your loved one is currently taking LIVMARLI or considering it, take advantage of this free program to connect with someone else like you. LIVMARLI Peers are patients or caregivers of children with Alagille syndrome who have experience with cholestatic pruritus and treatment with LIVMARLI.

You can discuss:

How they managed cholestatic pruritus in Alagille syndrome, what it was like to access LIVMARLI, and what their experience was with treatment with LIVMARLI.

Indication

LIVMARLI is a prescription medicine used to treat cholestatic pruritus (itch) in patients with Alagille syndrome 3 months of age and older. It is not known if LIVMARLI is safe and effective in children under 3 months of age or in adults 65 years and older.

Important Safety Information

What are the possible side effects of LIVMARLI?

LIVMARLI can cause serious side effects, including:

Please see next page for additional Important Safety Information.

Call 1-855-370-1371
or [CLICK HERE](#)
to register today!

livmarlipeerprogram.com

Important Safety Information Continued

Changes in liver tests. Changes in certain liver tests are common in patients with Alagille syndrome but may worsen during treatment with LIVMARLI.

These changes may be a sign of liver injury and can be serious. Your health care provider should do blood tests before starting and during treatment to check your liver function. Tell your health care provider right away if you get any signs or symptoms of liver problems, including:

- nausea or vomiting
- skin or the white part of the eye turns yellow
- dark or brown urine
- pain on the right side of the stomach (abdomen)
- loss of appetite

Stomach and intestinal (gastrointestinal) problems. LIVMARLI can cause stomach and intestinal problems, including diarrhea, stomach pain, and vomiting during treatment. Tell your health care provider right away if you have any of these symptoms more often or more severely than normal for you.

A condition called **Fat-Soluble Vitamin (FSV) Deficiency** caused by low levels of certain vitamins (vitamins A, D, E, and K) stored in body fat. FSV deficiency is common in patients with Alagille syndrome but may worsen during treatment. Your health care provider should do blood tests before starting and during treatment.

Other common side effects reported during treatment were gastrointestinal bleeding and bone fractures.

Tell your health care provider about all medicines that you take. LIVMARLI may affect the way some other medicines work, and some other medicines may affect the way LIVMARLI works. If you take a medicine that lowers cholesterol by binding bile acids, such as cholestyramine, colesevelam, or colestipol, take it at least 4 hours before or 4 hours after you take LIVMARLI. LIVMARLI is taken by mouth, 1 time each day, 30 minutes before a meal in the morning. Be sure to use the provided oral dosing dispenser to accurately measure the dose of medicine.

These are not all of the possible side effects of LIVMARLI. Call your health care provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

[Please click here for Patient Information](#)

